

P70532

Nomad Biosensors™ comprise a family of genetically encoded fluorescent sensors designed to monitor the signaling of G protein-coupled receptors (GPCRs) in cell-based assays.

Nomad Biosensors™ are engineered to measure the intracellular dynamics of second messengers such as calcium (Ca²⁺ Nomad), cAMP (cAMP Nomad), or diacylglycerol (DAG Nomad) upon GPCR activation. Additionally, β-arrestin signaling can also be studied using these biosensors. Nomad Biosensors™ can be combined in the same cell line for multiplex assays.

Prior to GPCR activation, the biosensors are localized in the plasma membrane. Upon ligand binding, the sensors undergo a conformational change that leads to an increase in fluorescence intensity and their relocalization within the vesicular trafficking pathways of the cells.



cAMPNOMAD DP1

cAMP Assay

Product Name: cAMPNomad-DP1 Cell Line

Reference: P70532

Gene Name: Prostaglandin D2 receptor (DP1)

cDNA Accession Number: NM000953

Host Cell Line: U2OS

Selection Markers: Geneticin (G418) + Hygromycin

Cell Quantity: > 3x10⁶ cells/vial


Storage Conditions: Liquid Nitrogen

About cAMPNomad-DP1


Nomad cell lines are a reliable system for studying G protein-coupled receptor (GPCR) signaling in living cells.

Optimized for the integration into High Content Screening (HCS) and High Throughput Screening (HTS) workflows, cAMPNomad-DP1 Cell Line stably express green cAMPNomad Biosensor along with the Prostaglandin D2 receptor (DP1).

Control



PGE2



cAMP Agonism Assay

The $cAMP$ Nomad-DP1 Cell Line was plated in a 96-well plate and incubated for a minimum of 4 hours and up to 24 hours at 37°C with 5% CO₂ to allow the cells to attach to the plate surface.

Agonism Assay: Cells were incubated with PGE2 diluted in a serum-reduced medium for 20–24 hours.

The increase in the fluorescence intensity of the green $cAMP$ Nomad biosensor (% Activity) was detected and analyzed using a microplate reader.




Figure 1. Dose-response curve for DP1 ligand.

Concentration response curve for PGE2 in the agonism assay.

The % Activity corresponds to the fluorescence intensity emitted by the green $cAMP$ Nomad biosensor normalized against the controls.