

P10970-IM


Chondrocytes are the resident cells of cartilage and are responsible for synthesizing a variety of collagenous and non-collagenous extracellular matrix macromolecules. These include collagen type II, aggrecan, and link protein, as well as smaller amounts of collagen types IX and XI. The regulation of chondrogenic cell proliferation and differentiation is central to the coordinated development of the vertebrate skeleton. Chondrocytes are capable of producing and responding to numerous peptide growth factors and cytokines, including insulin-like growth factor-1 and interleukin-1. Chondrocyte cultures serve as valuable in vitro models for studying cartilage regeneration and repair, the effects of cytokines and growth factors on cartilage, specific gene regulation, and the pathophysiology of arthritis.

Should you require further clarification or have specific inquiries, please do not hesitate to ask.



Innoprot

# IMMORTALIZED HUMAN CHONDROCYTES




**Product Type:** Immortalized Human Chondrocytes  
**Catalog Number:** IM- P10970  
**Immortalization:** SV40 Large T Antigen  
**Number of cells:** >5x10<sup>5</sup> cells (cryopreserved vials)  
**Storage:** Liquid Nitrogen  
**Recommended Medium:** Chondrocyte Medium (Reference: P60137)  
**Product Characterization:** Cytochemical staining for Collagen II, SB100B, CD56 and Aggrecan.




## About Immortalized Human Chondrocytes

The immortalized human chondrocyte cell line has been developed through the genetic modification of primary human chondrocyte cells, which are isolated from human articular cartilage. The SV40LT protein, derived from the Simian Virus 40 Large T-antigen, is used for this immortalization method. The introduction of the SV40LT protein confers immortality to the cells, significantly extending their lifespan compared to primary cells, which typically undergo senescence after a limited number of passages. Using SV40LT for immortalization is a common technique in cell biology, allowing for the establishment of cell lines with more stable and prolonged growth capacity. This modification enables researchers to conduct long-term experiments and studies requiring consistent cellular behavior over an extended period. While primary cells exhibit senescence after approximately five passages, the SV40LT-transduced cells demonstrate prolonged viability, extending beyond twenty passages.



Aggrecan CD56




Collagen II S100B

*This product contains a proprietary nucleic acid coding for a proprietary fluorescent protein intended for research purposes only. No rights are conveyed to modify or clone the gene encoding the fluorescent protein contained in this product or to use the gene or protein for any purpose other than non-commercial research, including compound validation or screening. For information on commercial licensing, contact the Licensing Department at Evrogen JSC via email: [license@evrogen.com](mailto:license@evrogen.com).*



## Alcian blue staining

One of the main characteristics of chondrocytes is their ability to synthesize a large amount of extracellular matrix. To analyze matrix synthesis and deposition, 40,000 cells/p24 were incubated for 12 days in complete medium in the absence or presence of ITS (insulin-transferrin-selenium). ITS promotes chondrogenesis and supports a differentiated phenotype. After 12 days, the cells were fixed with 4% PFA and stained with Alcian blue. Alcian blue staining, a cationic dye that stains sulfated glycosaminoglycans, confirmed an increase in glycan deposits in the presence of ITS.



## Culturing conditions

### 1. IMMEDIATELY UPON DELIVERY

- 1.1 Remove the vial from the shipping container to check for freezing.
- 1.2 Transfer the frozen vial to liquid nitrogen until ready to thaw.

### 2. THAWING CELLS

- 2.1 Prepare a poly-L-lysine coated flask (2  $\mu\text{g}/\text{cm}^2$ , T-75 flask recommended). Add 10 ml of sterile water to the T-75 flask, then add 150  $\mu\text{l}$  of poly-L-lysine stock solution (1 mg/ml, Innoprot Ref. PLL). Leave the flask in the incubator overnight or for a minimum of one hour at 37°C.
- 2.2 Prepare complete medium: Decontaminate the external surfaces of the medium and supplements with 70% ethanol, and transfer them to a sterile field. Aseptically open each supplement tube and add them to the basal medium with a pipette, rinsing each tube with the medium to recover the entire volume.
- 2.3 Rinse the poly-L-lysine coated flask twice with sterile water, then add 20 ml of complete medium to the flask. Leave the flask in the hood and proceed to thaw the cells.



## Culturing conditions

2.4 Place the vial in a 37°C water bath, holding and rotating it gently until the contents are completely thawed. Remove the vial immediately, wipe it dry, rinse with 70% ethanol, and transfer it to a sterile field. Remove the cap carefully without touching the interior threads. Gently resuspend the contents with a 1 ml Eppendorf pipette.

2.5 Dispense the vial contents into the equilibrated, poly-L-lysine coated culture vessels, recommending a seeding density higher than 6,500 cells/cm<sup>2</sup>.

*Note: Dilution and centrifugation of cells after thawing are not recommended as these actions can be more harmful than the effect of DMSO residue in the culture. It is also important to plate cells in poly-L-lysine coated culture vessels to promote cell attachment and growth.*

2.6 Replace the cap or cover of the flask, and gently rock the vessel to distribute the cells evenly. Loosen the cap if necessary to permit gas exchange.

2.7 Return the culture vessels to the incubator.

2.8 For best results, do not disturb the culture for at least 16 hours after initiation. Change the growth medium the next day to remove residual DMSO and unattached cells, then every other day thereafter.

### 3. MAINTENANCE OF THE CULTURE

3.1 Change the medium to fresh supplemented medium the morning after establishing the culture from cryopreserved cells.

3.2 Change the medium every three days thereafter until the culture is approximately 70% confluent.

3.3 Once the culture reaches 70% confluence, change the medium every other day until the culture is approximately 90% confluent.

### 4. SUBCULTURING

4.1 Subculture when the culture reaches 90% confluence or above.

4.2 Prepare poly-L-lysine-coated culture vessels (2 µg/cm<sup>2</sup>) one day before subculture.

4.3 Warm complete medium, trypsin/EDTA solution (T/E Solution), T/E neutralization solution (TNS), and DPBS (Ca<sup>++</sup>- and Mg<sup>++</sup>-free) to room temperature. We do not recommend warming reagents and medium in a 37°C water bath prior to use.

*Note: DPBS, trypsin/EDTA solution, and trypsin neutralization solution are included in the "Primary Cells Detach Kit" provided by Innoprot (Cat. N° P60305).*

4.4 Rinse the cells with DPBS.

4.5 Add 8 ml of DPBS and then 2 ml of T/E solution to the flask (for a T-75 flask). Gently rock the flask to ensure complete coverage of cells by T/E solution. Incubate at 37°C for 1 to 2 minutes or until cells completely round up. Use a microscope to monitor the change in cell morphology.

4.6 During incubation, prepare a 50 ml conical centrifuge tube with 5 ml of fetal bovine serum (FBS).

4.7 Transfer T/E solution from the flask to the 50 ml centrifuge tube (a small percentage of cells may detach) and continue incubating the flask at 37°C for another 1



## Culturing conditions

to 2 minutes (no solution should be in the flask at this time).

4.8 At the end of incubation, gently tap the side of the flask to dislodge cells from the surface. Check under a microscope to ensure all cells are detached.

4.9 Add 5 ml of TNS solution to the flask and transfer detached cells to the 50 ml centrifuge tube. Rinse the flask with another 5 ml of TNS to collect residual cells.

4.10 Examine the flask under a microscope to ensure a successful cell harvest, ensuring fewer than 5% of cells are left behind.

4.11 Centrifuge the 50 ml tube at 1000 rpm for 5 minutes. Resuspend cells in culture medium.

4.12 Count and plate cells in a new poly-L-lysine-coated culture vessel at the recommended cell density.



## Quality Control / Biosafety

The cells test negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi.

Caution: Handling human-derived products is potentially biohazardous. Although each cell strain tests negative for HIV, HBV, and HCV DNA, diagnostic tests are not necessarily 100% accurate. Proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working with these materials. Never mouth pipette. We recommend following universal procedures for handling products of human origin as the minimum precaution against contamination[1].

[1] Grizzle, W. E., & Polt, S. S. (1988). *Guidelines to avoid personal contamination by infective agents in research laboratories that use human tissues.* *J Tissue Culture Methods*, 11(4).

### **THIS PRODUCT IS FOR RESEARCH PURPOSES ONLY**

*It is not to be used for drug or diagnostic purposes, nor is it intended for human use. Innoprot products may not be resold, modified for resale, or used to manufacture commercial products without written approval of Innovative Technologies in Biological Systems, S.L.*